




Food and Agriculture
Organization of the
United Nations



Marine biodiversity of the Saya de Malha Bank shallows: A photographic catalogue



Marine biodiversity of the Saya de Malha Bank shallows: A photographic catalogue

by

Ranjeet Bhagooli

University of Mauritius, Moka, Mauritius

Sundy Ramah

Albion Fisheries Research Centre, Albion, Mauritius

Gilberte Gendron

Island Biodiversity and Conservation Centre,
University of Seychelles, and
Bee Ecological Consulting, Victoria

Deepeeka Kaullysing

University of Mauritius, Moka, Mauritius

Luvna Caussy

Albion Fisheries Research Centre, Albion, Mauritius

Edoardo Mostarda

Food and Agriculture Organization of the United Nations,
Rome

Required citation:

Bhagooli, R., Ramah, S., Gendron, G., Kaullysing, D., Caussy, L. & Mostarda, E. 2024. *Marine biodiversity of the Saya de Malha Bank shallows: A photographic catalogue*. FAO, Rome. <https://doi.org/10.4060/cd3735en>

The designations employed and the presentation of material in this information product do not imply the expression of any opinion whatsoever on the part of the Food and Agriculture Organization of the United Nations (FAO) concerning the legal or development status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. The mention of specific companies or products of manufacturers, whether or not these have been patented, does not imply that these have been endorsed or recommended by FAO in preference to others of a similar nature that are not mentioned.

The views expressed in this information product are those of the author(s) and do not necessarily reflect the views or policies of FAO.

ISBN 978-92-5-139479-3

© FAO, 2024



Some rights reserved. This work is made available under the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 IGO licence (CC BY-NC-SA 3.0 IGO; <https://creativecommons.org/licenses/by-nc-sa/3.0/igo/legalcode>).

Under the terms of this licence, this work may be copied, redistributed and adapted for non-commercial purposes, provided that the work is appropriately cited. In any use of this work, there should be no suggestion that FAO endorses any specific organization, products or services. The use of the FAO logo is not permitted. If the work is adapted, then it must be licensed under the same or equivalent Creative Commons licence. If a translation of this work is created, it must include the following disclaimer along with the required citation: "This translation was not created by the Food and Agriculture Organization of the United Nations (FAO). FAO is not responsible for the content or accuracy of this translation. The original [Language] edition shall be the authoritative edition."

Disputes arising under the licence that cannot be settled amicably will be resolved by mediation and arbitration as described in Article 8 of the licence except as otherwise provided herein. The applicable mediation rules will be the mediation rules of the World Intellectual Property Organization <http://www.wipo.int/amc/en/mediation/rules> and any arbitration will be conducted in accordance with the Arbitration Rules of the United Nations Commission on International Trade Law (UNCITRAL).

Third-party materials. Users wishing to reuse material from this work that is attributed to a third party, such as tables, figures or images, are responsible for determining whether permission is needed for that reuse and for obtaining permission from the copyright holder. The risk of claims resulting from infringement of any third-party-owned component in the work rests solely with the user.

Sales, rights and licensing. FAO information products are available on the FAO website (www.fao.org/publications) and can be purchased through publications-sales@fao.org. Requests for commercial use should be submitted via: www.fao.org/contact-us/licence-request. Queries regarding rights and licensing should be submitted to: copyright@fao.org.

All cover and interior photographs in this document: ©FAO/Norwegian Institute of Marine Research (IMR)

PREPARATION OF THIS DOCUMENT

The EAF-Nansen Programme “*Supporting the Application of the Ecosystem Approach to Fisheries Management considering Climate and Pollution Impacts*” of the Fisheries and Aquaculture Division of the Food and Agriculture Organization of the United Nations is an initiative to promote sustainable utilization of marine living resources and improved protection of the marine environment. To fulfill its objectives, the programme prioritizes the generation of scientific knowledge, with scientific surveys conducted aboard the research vessel (R/V) *Dr. Fridtjof Nansen*. Data are collected according to a science plan comprising 11 themes, with Theme 7 focusing on mapping critical and vulnerable habitats.

In May 2018, the programme conducted a survey on the Saya de Malha and Nazareth Banks with the objective of characterizing their marine ecosystems and morphology. Remotely Operated Vehicle dives were conducted in both shallow and deep waters of the area and subsequently, local scientists who had participated in the survey attended two workshops aimed at enhancing the identification of the benthic flora and fauna observed.

Following these workshops, a compilation of snapshots featuring key organisms observed during the dives and identified at the lowest taxonomic level possible was assembled. To optimize the use of these materials, it was suggested to compile these snapshots into a guide that would assist researchers in conducting future surveys in the region.

The EAF-Nansen Programme provided technical and financial support for the production of this catalogue. Moreover, the Joint Management Area, the mechanism of joint jurisdiction between Mauritius and Seychelles over the Saya de Malha Bank in the Mascarene Plateau region, endorsed the production of this catalogue to promote the sustainable management and understanding of the region’s marine life and biodiversity.

ABSTRACT

This photographic catalogue presents an overview of the flora and fauna observed in the shallow waters (less than 100 m depth) of the Saya de Malha Bank in the Mascarene Plateau. While not a comprehensive representation, as some taxa were excluded due to poor-quality photos that could result in misleading or unreliable depictions, it encompasses a total of 113 taxa identified at the family, genus, or species level. These taxa span key groups, including seaweeds, seagrasses, sponges, corals, echinoderms, crustaceans, molluscs, and fish.

The images are snapshots derived from video observations captured during the 2018 R/V *Dr. Fridtjof Nansen* ecosystem survey, which aimed to characterize the marine ecosystem and morphology of the Saya de Malha Bank. This area is notable for its rich marine biodiversity, supporting a diverse array of species and habitats that contribute to the overall health of oceanic environments.

This catalogue serves as a valuable resource for scientists conducting further explorations in the Mascarene Plateau region, providing a basic tool for taxonomic identification. By raising awareness of the significance of the Saya de Malha Bank, we aim to foster a greater appreciation for its unique marine life and promote the conservation of these critical habitats.

ACKNOWLEDGEMENTS

This work was made possible with the support of the EAF-Nansen Programme “*Supporting the Application of the Ecosystem Approach to Fisheries Management considering Climate Change and Pollution Impacts*” executed by Food and Agriculture Organization of the United Nations (FAO), with the scientific support of the Norwegian Institute of Marine Research (IMR), and funded by the Norwegian Agency for Development Cooperation (Norad). The authors are thankful to the FAO for funding and supporting Leg 2.1 of the regional resources and ecosystem surveys in the Indian Ocean (NORAD-FAO PROGRAMME GCP/GLO/690/NOR) on the Saya de Malha Bank on board the R/V *Dr Fridtjof Nansen* in 2018. The survey was carried out within the project framework entitled “*Characterising the marine ecosystem and morphology of the Saya de Malha Bank*” between the FAO and the Designated Authority (DA) of the Extended Continental Shelf, Mascarene Plateau region.

The authors are indebted to the Department for Continental Shelf, Maritime Zones Administration & Exploration (Prime Minister’s Office, Republic of Mauritius) for coordinating and co-leading the expedition and for their continuous support towards the publication of this photographic catalogue on the Marine Biodiversity of the Saya de Malha Bank Shallows, and the Mauritius-Seychelles Joint Commission of the Extended Continental Shelf for their support and assistance and granting the necessary authorisations.

During the R/V *Dr. Fridtjof Nansen* survey and two subsequent workshops, the authors of this photographic catalogue conducted identifications to the lowest taxonomic level possible within their areas of expertise across various groups. Subsequently, several experts were consulted and generously agreed to review the photographic material to enhance the accuracy of these preliminary identifications.

The authors of this catalogue extend their sincere gratitude to all these reviewers, whose assistance was crucial in validating or correcting the identifications. Their expertise greatly enhanced the presentation of a more accurate taxonomy in this catalogue.

Below, we list these experts by taxonomic group:

Sponges

Dr Michelle Kelly, National Institute of Water & Atmospheric Research, Auckland, New Zealand, and **Dr Toufiek Samaai**, Department of Forestry, Fisheries and the Environment, Cape Town, South Africa, emphasized that, as no specimens were examined, an element of doubt remains regarding the identity of these species based solely on images, particularly when they lack obvious diagnostic morphological characters that facilitate identification at the species level. The identifications provided here are the most accurate possible, based upon their personal understanding of the species in this and surrounding regions.

Corals

Scleractinian coral identifications were performed or verified by **Prof Francesca Benzoni**, King Abdullah University of Science and Technology, Saudi Arabia. In addition, other taxonomists provided their identifications, such as **Dr Lyndon Devantier**, Coral Reef Research, Townsville, Australia; **Dr David Obura**, CORDIO East Africa, Kenya; and **Dr Parisa Alidoost Salimi**, Department of Marine Biology, Science and Research Branch, Azad University, Tehran, Iran.

The authors compared the different identifications and, when they differed, decided to use the most conservative classification to ensure accuracy.

Echinoderms

Sea cucumber identifications were verified by **Prof Steve Purcell**, Marine Science and Management, National Marine Science Centre, Southern Cross University, Australia.

Starfish identifications were performed, improved, or verified by **Dr Christopher Mah**, Research Associate at the Smithsonian Museum of Natural History, USA.

Sea urchin identifications were performed or verified by **Dr Rich Mooi**, Curator of Echinoderms, Department of Invertebrate Zoology & Geology, California Academy of Sciences, USA.

Crustaceans

Prof Shane T. Ahyong, Senior Principal Research Scientist & Manager, Marine Invertebrates, Australian Museum Research Institute, Australia, identified the two species of stomatopods featured in this catalogue.

Bony fishes

Prof Kent Carpenter, Old Dominion University, USA, confirmed the identification of the two species of Lethrinids.

Dr Jeff Johnson, Manager of Ichthyology at the Queensland Museum, Australia, identified the sweetlip *Diagramma centurio*.

Dr Helen K. Larson, Curator Emeritus of Fishes at the Museum and Art Gallery of the Northern Territory, Australia, confirmed the identifications of the Gobiids.

Dr J. Howard Choat, James Cook University, Australia, confirmed the identification of the *Chlorurus* species.

Dr Benjamin Victor, Nova Southeastern University, USA, identified *Oxycheilinus bimaculatus*.

CONTENTS

ABOUT THE AUTHORS	xii
INTRODUCTION	1
The EAF-Nansen Programme	1
The Mauritius and Seychelles Joint Management Area	1
The Saya de Malha Bank.	3
Importance and uniqueness of the region	4
Biodiversity	4
The 2018 R/V <i>Dr Fridtjof Nansen</i> survey	5
Overview and objectives	5
Video transects data collection	6
Summary of key findings from the survey	6
Outputs	7
Purpose of the photographic catalogue and how to use it	7
SEAGRASSES AND SEaweEDS	9
SEAGRASSES	11
Family Cymodoceaceae	11
<i>Thalassodendron ciliatum</i>	11
Family Hydrocharitaceae	12
<i>Halophila decipiens</i>	12
SEaweEDS	13
Family Caulerpaceae	13
<i>Caulerpa cupressoides</i>	13
<i>Caulerpa taxifolia</i>	13
Family Halimedaceae	14
<i>Halimeda</i> sp. indet.	14
<i>Halimeda</i> sp. indet. 1.	14
Family Udoteaceae	15
<i>Udotea</i> sp. indet.	15
SPONGES	17
Family Agelasidae	19
<i>Agelas oxeata</i>	19
Family Axinellidae	19
<i>Axinella weltnerii</i>	19
<i>Dragmacidon coccineum</i>	20
Family Stelligeridae	20
<i>Higginsia pulcherrima</i>	20
Family Bubaridae	21
<i>Phakellia</i> sp. indet.	21
Family Scopalinidae	21
<i>Stylissa carteri</i>	21

Family Clionaidae	22
<i>Sphaciospongia inconstans</i>	22
Family Spongiidae	23
<i>Spongia</i> sp. indet.	23
Family Thorectidae	23
<i>Hyrtios erectus</i>	23
Family Chalinidae	24
<i>Haliclona</i> sp. indet.	24
Chalinidae gen. et sp. indet.	24
Family Petrosiidae	25
<i>Xestospongia testudinaria</i>	25
Family Hemiasterellidae	26
<i>Hemiasterella vasiformis</i>	26
<i>Liosina paradoxa</i>	26
Family Scleritodermidae.	27
<i>Aciculites tulearensis</i>	27

CORALS 29

HARD CORALS 31

Family Acroporidae	31
<i>Acropora</i> sp. indet.	31
<i>Acropora</i> sp. indet. 1	32
<i>Acropora</i> sp. indet. 2	32
<i>Acropora</i> sp. indet. 3	33
<i>Anacropora</i> sp. indet.	34
<i>Astreopora</i> sp. indet.	35
<i>Astreopora</i> sp. indet. 1.	35
<i>Montipora</i> cf. <i>undata</i>	36
Acroporidae gen. et sp. indet.	36
Family Agariciidae	37
<i>Leptoseris</i> sp. indet.	37
Family Fungiidae	38
Fungiidae sp. indet.	38
Fungiidae sp. indet. 1	38
Family Dendrophylliidae	39
<i>Turbinaria</i> cf. <i>mesenterina</i>	39
Family Lobophylliidae	40
<i>Lobophyllia</i> cf. <i>hemprichii</i>	40
Family Merulinidae	41
<i>Astrea</i> sp. indet.	41
<i>Dipsastraea</i> cf. <i>rotumana</i>	41
<i>Dipsastraea</i> sp. indet.	42
<i>Favites</i> cf. <i>abditata</i>	42
<i>Favites</i> cf. <i>vasta</i>	43
<i>Oulophyllia crispera</i>	43
<i>Paramontastraea peresi</i>	44

<i>Platygyra</i> cf. <i>acuta</i>	44
<i>Platygyra</i> cf. <i>daedalea</i>	45
Family Pocilloporidae	45
<i>Pocillopora</i> sp. indet.	45
<i>Pocillopora</i> sp. indet. 1.	46
<i>Stylophora</i> sp. indet.	46
Family Poritidae	47
<i>Goniopora</i> sp. indet.	47
<i>Porites</i> sp. indet.	48
<i>Porites</i> sp. indet. 1 (Growth form I)	49
<i>Porites</i> sp. indet. 2 (Growth form II)	50
OCTOCORALS	51
Family Helioporidae	51
<i>Heliopora</i> cf. <i>coerulea</i>	51
<i>Heliopora</i> sp. indet.	52
Family Ellisellidae	53
Ellisellidae gen. et sp. indet.	53
Family Isididae	53
Isididae gen. et sp. indet.	53
<i>Rumphella</i> sp. indet.	54
Family Sarcophytidae.	55
<i>Lobophytum</i> sp. indet.	55
<i>Sarcophyton</i> sp. indet.	55
HYDROZOANS	56
Family Milleporidae	56
<i>Millepora</i> sp. indet.	56
ECHINODERMS	57
SEA CUCUMBERS	59
Family Holothuriidae	59
<i>Holothuria atra</i>	59
<i>Holothuria coronopertusa</i>	59
Family Stichopodidae.	60
<i>Stichopus</i> cf. <i>pseudohorrens</i>	60
<i>Thelenota ananas</i>	60
STARFISHES	61
Family Ophidiasteridae	61
<i>Linckia</i> cf. <i>laevigata</i>	61
<i>Linckia</i> sp. indet.	61
Family Oreasteridae	62
<i>Culcita</i> cf. <i>novaeguineae</i>	62
<i>Culcita schmideliana</i>	62
Family Ophidiasteridae	62
<i>Pentaceraster</i> sp. indet.	63
<i>Protoreaster lincki</i>	63

SEA URCHINS	64
Family Diadematidae	64
<i>Echinothrix calamaris</i>	64
<i>Echinothrix diadema</i>	64
Family Cidaridae	65
Cidaridae gen. et sp. indet.	65
Family Toxopneustidae	65
<i>Toxopneustes pileosus</i>	65
OTHER INVERTEBRATES	67
MOLLUSCS	69
Family Conidae	69
<i>Conus</i> cf. <i>tessulatus</i>	69
Family Strombidae	69
<i>Lambis</i> cf. <i>lambis</i>	69
Family Phyllidiidae	70
<i>Phyllidiopsis quadrilineata</i>	70
CRUSTACEANS	71
Family Odontodactylidae	71
<i>Odontodactylus japonicus</i>	71
<i>Odontodactylus latirostris</i>	71
BONY AND CARTILAGINOUS FISHES	73
BONY FISHES	75
Family Muraenidae	75
<i>Gymnothorax favagineus</i>	75
Family Holocentridae	75
<i>Myripristis botche</i>	75
<i>Myripristis</i> sp. indet.	76
<i>Sargocentron</i> cf. <i>seychellense</i>	76
<i>Sargocentron</i> cf. <i>diadema</i>	77
Family Apogonidae	77
<i>Apogonichthyoides pharaonis</i>	77
Family Gobiidae	78
<i>Amblyeleotris periophthalma</i>	78
<i>Valenciennesa</i> cf. <i>wardii</i>	79
Family Trichonotidae	79
<i>Trichonotus</i> cf. <i>marleyi</i>	79
Family Callionymidae	80
<i>Synchiropus</i> cf. <i>sechellensis</i>	80
Family Fistulariidae	80
<i>Fistularia</i> cf. <i>commersonii</i>	80
Family Carangidae	81
Carangidae gen. et sp. indet.	81
Family Echeneidae	81
<i>Echeneis naucrates</i>	81

Family Epinephelidae	82
<i>Variola louti</i>	82
Family Scorpaenidae	82
<i>Dendrochirus hemprichi</i>	82
<i>Pterois miles</i>	83
Family Zanclidae	83
<i>Zanclus cornutus</i>	83
Family Acanthuridae	84
<i>Naso brevirostris</i>	84
<i>Naso hexacanthus</i>	84
Family Haemulidae	85
<i>Diagramma centurio</i>	85
Family Lutjanidae	85
<i>Lutjanus sebae</i>	85
Family Lethrinidae.	86
<i>Lethrinus enigmaticus</i>	86
<i>Lethrinus variegatus</i>	87
Family Nemipteridae	87
<i>Scolopsis frenata</i>	87
Family Pomacanthidae.	88
<i>Pomacanthus imperator</i>	88
<i>Pomacanthus semicirculatus</i>	88
Family Labridae.	89
<i>Oxycheilinus bimaculatus</i>	89
<i>Chlorurus</i> sp. indet.. . . .	89
Family Balistidae.	90
<i>Sufflamen chrysopterum</i>	90
Family Ostraciidae.	90
<i>Lactoria fornasini</i>	90
Family Tetraodontidae	91
<i>Arothron hispidus</i>	91
<i>Canthigaster rivulata</i>	91
CARTILAGINOUS FISHES	92
Family Dasyatidae.	92
<i>Taeniurops meyeri</i>	92
REFERENCES	93
FURTHER READINGS	94
INDEX	95

ABOUT THE AUTHORS

Dr Ranjeet Bhagooli is a Professor in Marine Biology & Biological Oceanography at the Department of Biosciences and Ocean Studies, Faculty of Science, University of Mauritius (UoM), an Adjunct Professor at Diponegoro University, Indonesia and an Associate Fellow at the Institute of Oceanography and Environment, Universiti Malaysia Terengganu, Malaysia. He holds an MSc in Biology, Chemistry and Marine Science, and a PhD in Marine and Environmental Sciences from the University of Ryukyus, Okinawa, Japan. He was a post-doctoral Research Associate and Fellow at the University of Technology, Sydney / University of Tasmania, Australia and University of Miami, Rosenstiel School of Marine, Atmospheric and Earth Science, Florida, USA, respectively. He leads the Pole of Research Excellence in Sustainable Marine Biodiversity at UoM.

Dr Sundy Ramah is a Scientific Officer at the Albion Fisheries Research Centre under the Ministry of Agro-Industry, Food Security, Blue Economy and Fisheries (Blue Economy and Fisheries Division) in Mauritius. He holds a BSc (Hons) in Marine Science and Technology and a PhD in the field of Marine Ecophysiology & Molecular Biology from the University of Mauritius. He is a member of the Western Indian Ocean Marine Science Association (WIOMSA) and a professional member of the Marine Biological Association (UK).

Gilberte Gendron is a marine biologist with extensive expertise in marine biodiversity and ecosystem conservation. She holds an MSc in Oceanography from the Université du Québec à Rimouski, Canada. She is the Director of Bee Ecological Consulting and a core member of the Island Biodiversity & Conservation (IBC) Center at the University of Seychelles. With a strong background in marine research and applied environmental consulting, she has contributed significantly to strategic conservation initiatives across in Seychelles and the Western Indian Ocean region.

Dr Deepeeka Kaullysing is a Senior Lecturer at the Department of Biosciences and Ocean Studies, Faculty of Science, University of Mauritius, Republic of Mauritius. She holds an MSc in Marine Science with specialization in Marine Biology from Goa University and a PhD in Marine Malacology and Ecology from the University of Mauritius. She is a member of the Pole of Research Excellence in Sustainable Marine Biodiversity at the University of Mauritius.

Dr (Ms) Luvna Caussy is a Scientific Officer at the Albion Fisheries Research Centre under the Ministry of Agro-Industry, Food Security, Blue Economy and Fisheries (Blue Economy and Fisheries Division) in Mauritius. She holds a BSc (Hons) in Agriculture with specialisation in Aquaculture and a PhD Fisheries Biology from the University of Mauritius.

Dr Edoardo Mostarda is a fisheries and fish biodiversity expert at the Food and Agriculture Organization of the United Nations (FAO) in Rome, Italy. After a decade of work as a fish ecologist, he joined the FAO FishFinder Programme (formerly the Species Identification and Data Programme) in 2011. Since then, he has contributed to the development of over 25 species identification tools, serving as an author, technical editor, or scientific editor in various FAO projects, including the EAF-Nansen Programme.

INTRODUCTION

The EAF-Nansen Programme

The EAF-Nansen Programme is a longstanding partnership between the Food and Agriculture Organization of the United Nations (FAO) and Norway, in close collaboration with the Norwegian Institute of Marine Research (IMR). Working with 32 countries across Africa and the Bay of Bengal, the programme aims to enhance food and nutrition security by strengthening fisheries policies and management practices in line with the ecosystem approach to fisheries (EAF).

The programme is structured around three interconnected areas that contribute to sustainable fisheries and improved food and nutrition security for people in partner countries: scientific advice, fisheries management, and capacity development. Scientific knowledge is generated through surveys conducted aboard the research vessel *Dr. Fridtjof Nansen*, the primary tool for collecting fisheries-independent data. The vessel's work, through the collection of unique data, contributes to monitoring and assessing fisheries resources while addressing pressing global concerns such as biodiversity conservation, and the impacts of climate change, pollution, and human activities on marine ecosystems. The programme's science plan guides the efficient collection and use of data to support research and decision-making.

Theme 7 of the science plan aims to provide comprehensive information on bottom habitats, particularly focusing on the vulnerable ones. These habitats, which are at risk from fisheries and other human activities, often include biodiversity hotspots that recover slowly from disturbances and thus qualify as vulnerable marine ecosystems (VMEs). Understanding their characteristics and distribution is crucial for providing science-based advice for targeted and effective protective measures. The programme's bottom habitat studies are primarily focused on the deep-sea areas beyond national jurisdiction (ABNJ), where there is still limited knowledge about species and habitat diversity. Studies within national waters are also conducted, especially in relation to oil and gas activities, to provide baseline data for environmental impact assessments of such projects.

This photographic catalogue was developed to support the work of Theme 7 and scientists conducting further explorations in the Mascarene Plateau region, providing a tool for accurate taxonomic identification and enhancing future research and conservation efforts.

The Mauritius and Seychelles Joint Management Area

In March 2011, the United Nations Commission on the Limits of the Continental Shelf (CLCS) adopted recommendations confirming the entitlement of the Republic of Mauritius and the Republic of Seychelles to an area of extended continental shelf in the

Mascarene Plateau region, as detailed in a joint submission by the two states. In March 2012, Mauritius and Seychelles signed two treaties. The first treaty established the outer limits of the extended continental shelf and outlined their joint sovereign rights for exploration and resource exploitation, in accordance with the 1982 United Nations Convention on the Law of the Sea (UNCLOS). The second treaty, effective for 30 years, created the Joint Management Area (JMA), which spans approximately 396 000 km² and is jointly managed by both countries (Figure 1).




Figure 1 Map of the Joint Management Area (JMA) over the Continental Shelf in the Mascarene Plateau. **Source:** Author's elaboration on background grid: GEBCO Compilation Group (2024) GEBCO 2024 Grid (doi:10.5285/1c44ce99-0a0d-5f4f-e063-7086abc0ea0f). Conforms to UN Map of the World, 2024.

This treaty is particularly significant as it represents the first instance of joint management over such a large maritime area and establishes an institutional and regulatory framework for the sustainable management and exploitation of its natural resources. Both Mauritius and Seychelles are committed to the 2030 Agenda for Sustainable Development and the Paris Agreement on climate change, with a strong emphasis on Sustainable Development Goal (SDG) 14, which focuses on marine life. They reaffirm their dedication to sustainably managing the JMA's natural resources, prioritising the precautionary principle and the protection of the marine environment and biodiversity.

The Saya de Malha Bank

A notable feature within the JMA is the Saya de Malha Bank, a vast submerged feature extending between Mauritius and Seychelles, spanning roughly from 9° to 13° S and 59° to 63° E (Figure 2).




Figure 2 Map of the Saya de Malha Bank. The red dots are the stations where the VAMS was deployed. **Source:** Author’s elaboration on background grid: GEBCO Compilation Group (2024) GEBCO 2024 Grid (doi:10.5285/1c44ce99-0a0d-5f4f-e063-7086abc0ea0f). Conforms to UN Map of the World, 2024

It is among the most remote and least explored shallow water ecosystems globally with the closest land, Agalega, approximately 300 km to the west. It is also among the world's largest underwater Banks; the smaller North Bank, also known as the Ritchie Bank, and the significantly larger and more southerly Saya de Malha Bank together cover an area of 40 808 km², largely within international waters. The depths on these Banks vary from 8 to 150 m, separated by a 1 000 m deep channel. The Banks gradually slope downwards on all sides into the deep open ocean. Ridges and pinnacles are scattered on the Bank's relatively flat, plateau-like topography, creating a complex underwater landscape. The Saya de Malha Bank is influenced by the South Equatorial Current (SEC) which enhances oceanic productivity in the region. The seabed of Saya de Malha Bank is jointly managed by the Republics of Mauritius and Seychelles, while the water column remains in the high seas.

Importance and uniqueness of the region

Scientific research on the underlying seafloor of the Saya de Malha Bank has become primordial to help both the Republic of Mauritius and the Republic of Seychelles in the future development, management, and effective conservation strategies within the JMA using science-based evidence. The Saya de Malha Bank has been identified as globally important in two independent processes (Obura *et al.*, 2012):

- the technical process supporting the Convention on Biological Diversity used the Bank as an illustration of a likely Ecologically or Biologically Significant Area (EBSA), satisfying four of seven criteria: 1. Uniqueness or rarity; 2. Special importance for life history stages of species; 3. Importance for threatened, endangered, or declining species and/or habitats; and 5. Biological productivity; and
- a separate analysis by World Wildlife Foundation, as part of the Western Indian Ocean Marine Ecoregion (WIOMER) and 'Réseau des Aires Marines Protégés' project of the Indian Ocean Commission (RAMP-COI) scored Saya de Malha as high as the other 3 EBSA criteria, namely: 4. Vulnerability, Fragility, Sensitivity, or Slow recovery; 6. Biological diversity; and 7. Naturalness; and has selected it as a priority seascape of global significance for its eco-regional conservation strategy.

Biodiversity

The Saya de Malha Bank is a biodiversity hotspot thanks to oceanic productivity, enhanced by the interaction of its topography and the SEC. In addition to seagrasses, the Bank hosts extensive coral communities and encrusting red coralline algae, while more than 150 species of invertebrates and 100 species of gastropods (a diverse class of molluscs) have been identified (Hilbertz *et al.*, 2002). Among the array of fauna are parrotfish, surgeonfish, rabbitfish, green sea turtles, wedge-tailed shearwaters, white-

tailed tropicbirds, spotted dolphins, pygmy blue whales, and pilot whales. The coral reefs of Saya de Malha are a critical stepping stone for the migration of shallow-water species across the Indian Ocean, and sailors crossing the Banks have reported that they are also a major breeding ground for sperm whales and blue whales (Webber *et al.* 2023).

Current knowledge holds that Saya de Malha Bank hosts the most abundant green calcareous algae (*Halimeda* spp.) field and probably constitutes the main source of unstable substrate (sand) dominating the stations surveyed. Thirteen genera of corals have previously been recorded from the shallow areas of the SMB (Rosen, 1971). At the north Bank, however, an entirely different coral ecosystem was found, dominated by large stands of a single species of branching *Acropora* corals (Hagan and Robinson, 2001). *Acropora*, *Pocillopora*, *Montipora*, *Porites*, *Heliopora*, *Danafungia*, *Favites*, *Dipsastraea*, *Seriatopora* and the possibility of *Stylophora* were the most common genera amongst others.

The 2018 R/V *Dr Fridtjof Nansen* survey

Overview and objectives

The cruise entitled “Characterizing the Marine Ecosystem and Morphology of the Saya de Malha Bank” started on 3 May 2018 when the Research Vessel *Dr Fridtjof Nansen* left Port Victoria, Seychelles, and ended on 4 June 2018 when the vessel docked in Port Louis, Mauritius.

The overriding general aim of the cruise was to characterize the marine ecosystem and morphology of the Saya de Malha Bank. Sampling using the full complement of technologies of the vessel facilitated investigations of the following aspects:

- geomorphology, benthic habitats, and benthos (Multibeam sounder mapping in subareas and along pre-determined transects, habitat and benthos studies emphasising sandy subareas and subareas with macroalgae, seagrass, and coral);
- fish and crustacean resources (density mapping with acoustic and optical technologies, emphasising commercial and toxic species); and
- physical and chemical oceanography, including current measurements (onboard measurements).

In addition, sampling in support of studies with a wider geographical scope undertaken on other surveys under the EAF-Nansen Programme were conducted, e.g. tissue sampling for genetics, contaminants, mammal observations, recording of microplastics and litter.

One key question the cruise aimed at addressing is in regard to geomorphology, benthic habitats, and benthos. Coarse maps of the bathymetry and substrates were generated from previous surveys. The technologies available on the research vessel

facilitate more detailed and accurate mapping that will be useful as baselines for further scientific studies and assessments related to industrial developments. This was done using single- and multibeam acoustics and the sub-bottom profiler to map geomorphology in selected subareas and along pre-determined transects, and use the video-assisted multisampler (VAMS) to collect grab samples and HD videos for benthos studies. Benthos studies were focused on soft-bottom substrates and along upper slopes where coral cover is expected. Data gathered from the VAMS videos recorded in shallow areas of Saya de Mahla were used to develop this catalogue.

Video transects data collection

In-situ visual data collection was carried out using the HD camera of a tethered Remotely Operated Vehicle (ROV), able to operate between 20 and 1000 m depth, attached to a Video-assisted Multi-sampler (VAMS), a tubular cage hosting the ROV (Serigstad *et al.*, 2015).

The ROV attached to the VAMS was used to carry out the video transect survey in two modes: 1) as a point sampler where the VAMS was deployed on the seabed allowing the ROV to explore the immediate 15 m vicinity in four directions (north, east, south and west) and 2) in a towed mode whereby the vessel towed the VAMS at 5 m above the seabed along pre-determined transects perpendicular to isobaths at 0.1–0.4 knots while the ROV explored the underlying seabed. The distance between the ROV camera and the seabed was kept at 2 m, lens' swath width of 15 m, during the steady passage, depending on the environment and sea condition.

As a first exploratory survey, several stops were made along the transects in order to have close-up videos for record and identification of certain benthic organisms. The video records were logged with data on GPS position, to calculate the distance covered by the ROV along the transect, dive time and depth in the CAMPOD Logger software, a camera-based data collection software designed by the Institute of Marine Research of Norway (IMR) used with the ROV.

The CAMPOD Logger uses the same principle as the Coral Point Count with Excel extension (CPCe) Software (Kohler and Gill, 2006). The data was further analysed after the cruise.

Summary of key findings from the survey

For the benthos, the shallow water flats (depth 20–70 m) were dominated by organisms such as seagrass (*Thalassodendron ciliatum*), calcareous red algae (*Lithothamnion* sp.), and green algae (*Halimeda* sp.). In various locations where *Halimeda* species were dominant, the upper 5 cm layer of sediment consisted mainly of its dead calcareous leaves. Shallow waters also provided a range of degraded coral colonies, including branching *Porites*, *Acropora* sp., *Heliopora* sp., *Tubastrea* sp., *Cyphastrea* sp., *Turbinaria* sp., *Fungia* sp., *Favites* sp., *Favia* sp., and *Galaxea* sp., which were more dominant than others. These calcified algae and corals, together with foraminiferans, are likely to be important in forming reef construction and constitute a notable portion of the bottom sediments—sand and gravel—in the investigated area. An additional source of bottom sediments and reefal boulders comes from remnants of fossil gastropods and bivalves.

The areas predominantly occupied by the organisms listed above were well recognized during visual observations: *Thalassodendron ciliatum*, *Halimeda* sp., and corals (with *Lithothamnion*) or only *Lithothamnion*. Occasionally, sands with no epibenthic domination were observed. These five communities likely represent different stages of succession.

The deeper locations were divided into two zones: 40–100 m and 100–1000 m. The first zone was characterized by very poor fauna, with flat, sandy rocky slopes observed. In the 100–1000 m depth zone, a more diverse fauna was observed on different substrates, which were mainly dominated by sand or rocks.

Additional findings related to physical oceanography, geomorphology and benthic substrates, chemical oceanography and phytoplankton, zooplankton, pelagic and demersal fish, mammals, birds, and turtles, as well as other observations (including litter and microplastics), can be viewed in the outputs described below.

Outputs

This photographic catalogue is among one of the outputs of the 2018 Nansen Research Cruise. The details of the cruise, including activities and planned outcomes and outputs, have been documented mainly in the NORAD–FAO project GCP/INT/730/NOR cruise reports “*Dr. Fridtjof Nansen – Regional Resources and Ecosystem Survey in the Indian Ocean, Leg 2.1. Characterizing ecosystems and morphology of the Saya de Malha Bank and Nazareth Bank, Survey no 2018406, 3 May–4 June, 2018, by Bergstad et al. (2018).*

The scientific papers emanating from the surveys and follow-up analyses during the post-cruise workshops have been published in a Special Issue edition of the Western Indian Ocean Journal of Marine Science (WIOJMS) (<https://www.wiomsa.org/publications-2/wio-journal-of-marine-science/>) hosted by the Western Indian Ocean Marine Science Association (WIOMSA) (<https://www.wiomsa.org/>). The special edition was entitled “Studies on the Mascarene Plateau” and the main findings were presented under topics like ocean circulation patterns; inorganic nutrient distribution; chlorophyll a and microphytoplankton density and diversity distribution; shallow-water benthic habitat diversity and distribution; sediment grain size and organic matter contents; photosynthetic functioning of macroalgae, seagrass, healthy and bleached scleractinian corals; marine molluscs diversity; macro- and mega-fauna of the Bank slopes; pelagic and demersal fish diversity; field observations on *Halimeda*, Rhodoliths and *Thalassodendron* beds, the coral-eating Crown-of-Thorn and other starfish, and endemic giant clam sightings; and coral disease.

The full text articles are accessible at <https://www.ajol.info/index.php/wiojms/issue/view/20888>

Purpose of the photographic catalogue and how to use it

This photographic catalogue is designed to assist in identifying a selection of taxa observed during the video surveys of the 2018 R/V *Dr Fridtjof Nansen* survey.

It is not meant to be a comprehensive representation of the fauna and flora of the

shallow-water habitats of the Saya de Malha Bank. Several taxa were observed and tentatively identified; however, they were excluded from this catalogue due to the poor quality of the photos, which would have resulted in a misleading or unreliable representation. The photos and descriptions provided are based on the author's expertise and that of numerous reviewers. However, identifying marine species can be complex, and while every effort has been made to ensure accuracy, there are inherent limitations to this catalogue.

Many marine organisms exhibit variations in colour, size, and behaviour that can make identification challenging. The taxa included here have been identified to the best of the authors' and reviewers' knowledge, but they may not account for all variations or look-alikes. We recommend consulting additional resources or seeking expert advice when precise identification is critical.

The taxa included in this catalogue are organized by major taxonomic groups: seagrasses and seaweeds, sponges, corals, echinoderms, other invertebrates, and bony and cartilaginous fishes. Within each group, taxa are arranged alphabetically by class, order, and family.

For each taxon, the scientific name is provided, including authorship. If the species could not be identified, the genus name is followed by "sp. indet." If both genus and species are undetermined, the family name is followed by "gen. et sp. indet."

A brief description highlights key characteristics used for identification, along with the habitat where the taxon was observed. Each photograph is labeled with the station where it was captured and the depth in metres. The locations of the stations are shown in Figure 2.

SEAGRASSES AND SEAWEEDS



Thalassodendron ciliatum (Forsskål) Hartog, 1970



St. 13. Depth: 31 m



St. 36. Depth: 28 m

A common seagrass species, forming dense meadows. It has long, ribbon-like, dark-green leaves that are linear and often have serrated edges, resembling blades or straps. The plant's overall shape can be described as bushy or tufted, with leaves radiating from a central point or growing in clusters along rhizomes.

Habitat: sandy or muddy substrates.

Halophila decipiens Ostenfeld, 1902



St. 34. Depth: 45 m



St. 34. Depth: 45 m

A small seagrass species, with elongated, paddle-shaped leaves arranged in pairs along the stem. The smooth leaves range from bright green to olive-green. **Habitat:** shallow waters, on sandy or muddy substrates with clear water and moderate wave action.

Caulerpa cupressoides (Vahl) C.Agardh, 1817



A distinctive seaweed with upright branches resembling miniature trees or cacti. The branches are arranged in a dense, bushy manner, giving the alga a compact and sculptural form. **Habitat:** rocky substrates or coral reefs.

Caulerpa taxifolia (Vahl) C.Agardh, 1817



A fast-growing seaweed with upright, feather-like branches, standing 3 to 10 cm tall. Each leafy branch is flat, slightly curved upwards, and tapers at both ends. Colour varies from dark to light green. **Habitat:** sandy bottoms, rocky outcroppings, mud, and natural meadows.


Halimeda sp. indet.



St. 36. Depth: 38 m

A frequently encountered seaweed species characterized by distinctive connected calcified segments, reminiscent of a chain of coins. In this species, the segments are round and arranged in flat, fan-like structures. **Habitat:** shallow coral reefs.

Halimeda sp. indet. 1



St. 34. Depth: 46 m



A different species of *Halimeda* characterized by a branched structure and oval segments, and found in isolated patches or scattered individuals. **Habitat:** sandy bottoms, attached to dead corals.

Udotea sp. indet.



A seaweed species with a fan-shaped thallus with flattened, branching blades with calcified segments. It is anchored to the bottom by uncalcified tufts of rhizoids, which vary in shape depending on the substrate. **Habitat:** *Udotea* species grow isolated or in groups and prefer shallow, well-lit habitats such as coral reefs, where they contribute to the benthic ecosystem's structure and function.

SPONGES



Family Agelasidae

Class Demospongiae
Order Agelasida

Agelas oxeata Lévi, 1961



St. 39. Depth: 41 m

A lobed-shaped sponge with a velvety surface texture that contributes to its distinctive appearance. The sponge is characterized by a brownish-red hue, which is a key diagnostic feature for identifying the species in its natural habitat.

Habitat: coral reefs and rocky habitats.

Family Axinellidae

Order Axinellida

Axinella weltnerii (Lendenfeld, 1897)




St. 37. Depth: 33 m

An erect, flabellate or fan-shaped sponge with a rough, twisted, and wrinkled surface with shallow ridges. Colour is a dark red. **Habitat:** rocky or sand rubble substrates.

Family Axinellidae

Class Demospongiae
Order Axinellida

Dragmacidon coccineum (Keller, 1891)




St. 42. Depth: 49 m

This sponge has an irregularly encrusting form with a smooth surface that ranges from faintly domed to undulating. Oscules are regularly spaced and slightly elevated. It is brilliant red in color, with a whitish sheen caused by surface sand granules. **Habitat:** found on coral reefs.

Family Stelligeridae

Higginsia pulcherrima Pulitzer-Finali, 1993



St. 4. Depth: 33 m

This sponge features an irregularly contoured structure with multiple blades extending from a common base. Externally, it displays a dull orange coloration with ridged-shaped blades. **Habitat:** found on coral rubble.

Family Bubaridae

Class Demospongiae
Order Bubarida

Phakellia sp. indet.



An undetermined species of *Phakellia*, which includes 36 species worldwide. *Phakellia* species are generally fan-shaped with thin plates. Its fans are reinforced by thick spicule tracts, resembling veins or stout lines. **Habitat:** rocky flats and seagrass/seaweed beds.

Family Scopalinidae

Order Scopalinida

Stylissa carteri (Dendy, 1889)



An erected, lamellar sponge, usually fan-shaped, with short, stout projections covering both surfaces, forming extensive ridges that diverge towards the edges. Bright orange in colour. **Habitat:** rocky substrates.

Sphaciospongia inconstans (Dendy, 1887)



St. 34. Depth: 44 m



St. 40. Depth: 45 m

A massive sponge exhibiting an irregularly sub-spherical shape, with a shallow, slightly raised depression at its summit housing numerous oscules. Its surface presents a uniform appearance, devoid of protruding structures, yet adorned with numerous distinctive circular or meandering openings seamlessly integrated with the surface. Colour brownish to golden. **Habitat:** reef flats and seagrass beds.

Spongia sp. indet.



An undetermined species of *Spongia*, which includes 78 valid species worldwide. Members of this genus are soft to firm and compressible, with a skeletal structure that includes primary fibers with foreign inclusions, and a meshwork of interconnecting secondary fibers providing flexibility and water retention. **Habitat:** sandy bottoms.

Family Thorectidae

Hyrtios erectus (Keller, 1889)



A characteristic sponge forming a club-shaped lobe that extends vertically from the substrate and features a rounded apex with the oscules, located at the apex. Its overall coloration is brown-black. **Habitat:** reefs and rocky substrates.

Haliclona sp. indet.



This species belongs to the Chalinid sponges and is assigned to the genus *Haliclona*. The genus *Haliclona* comprises approximately 480 valid species globally, exhibiting a wide range of sponge forms, from erect and tube-shaped to branching structures. The textures of these sponges can vary significantly, from soft and fragile to firm and elastic. **Habitat:** *Haliclona* sponges are found across various depths but are more commonly observed in relatively shallow environments.

Chalinidae gen. et sp. indet.



A Chalinid sponge which could not be assigned to one of the five genera in the family. Taxonomic research within this family presents challenges due to the scarcity, simplicity, and occasional variability in distinguishing characteristics among its species.

Xestospongia testudinaria (Lamarck, 1815)



Commonly referred to as the giant barrel sponge, this species is morphologically unique with its large, barrel-shaped form and tough, rigid exterior. It displays a maroon to pink coloration, with a pale white opening at the top. These sponges can grow to notable sizes, reaching up to 1.5 m in height and 1 m in diameter. **Habitat:** this species primarily inhabits shallow tropical waters, especially within coral reef environments. It is commonly found in areas with hard substrates that allow it to anchor securely. It can also be found in deeper waters, extending to depths of up to 55 m. In the Saya de Malha Bank, most specimens were observed in *Halimeda* meadows.


Hemiasterella vasiformis (Kirkpatrick, 1903)



St. 42. Depth: 49 m

A large, typically cup-shaped sponge, characterized by a slightly thickened base, firm and rubbery texture, and pinkish-purple coloration. **Habitat:** found on rocky or sand rubble substrates.

Liosina paradoxa Thiele, 1899



St. 37. Depth: 37 m

A sponge that grows in a massive, lobose form, rising vertically from a broad base. It features branches extending upward, with oscules located at the apex. The surface appears tuberculated, with a muted, muddy grey coloration. **Habitat:** typically found on rocky substrates.

Aciculites tulearensis Vacelet & Vasseur, 1965



St. 34. Depth: 45 m

This species is a relatively large sponge with an irregularly contoured, thick blade shape. Its inhalant side features visible subdermal canals, while the exhalant side is marked by small conical protrusions with oscules. Externally, the sponge appears dull dark brown, while internally it is a paler brown. **Habitat:** initially reported from dark tunnels and cavities within barrier reefs. It predominantly inhabits rocky substrates and is commonly associated with coral reefs.

CORALS



Acropora sp. indet.



Colonies exhibit an arborescent growth pattern, characterized by cylindrical branches that intertwine to form thickets. The axial corallites (the stony cups in which the polyps sit) are prominently exerted. These colonies typically display a cream to brown coloration, with pale branch ends. **Habitat:** commonly found inhabiting both the reef slopes and flats.

***Acropora* sp. indet. 1**





Colonies consist of open thickets of sturdy, tapered branches. Axial corallites are moderately large and tubular to dome-shaped. Radial corallites are of mixed sizes, widely spaced, and vary in length. The colonies are brown with pale tips.
Habitat: observed in shallow reef environments.

***Acropora* sp. indet. 2**




Colonies exhibit a digitate morphology, characterized by finger-like branches arising from an encrusting, plate-like base. Branches are thick, gradually tapering to large, dome-shaped axial corallites. The colonies are brown with cream tips.
Habitat: observed on exposed upper reef slopes and reef flats.

Acropora sp. indet. 3



Colonies are arborescent with cylindrical branches and horizontally fused basal branches that have upturned ends. Radial corallites are of mixed sizes. The colonies are pale to dark brown with paler tips. **Habitat:** restricted to shallow back-reef margins.

Anacropora sp. indet.




St. 39. Depth: 43 m



St. 39. Depth: 43 m


A species of *Anacropora* with evenly spaced branches, thick and tapered, with blunt tips. The coenosteum, the skeletal material lying between the corallites, is smooth. Colonies are pale brown with white branch tips. **Habitat:** observed in reef environments.

***Astreopora* sp. indet.**




Colonies are hemispherical, with an even surface. Corallites are evenly spaced and conical with rounded, upright or outwardly directed openings. The coenosteum has outwardly directed papillae. They are brownish in colour. **Habitat:** observed in reef environments.

***Astreopora* sp. indet. 1**



Colonies take on an encrusting form, with thick-walled and conical corallites. The coenosteum between the corallites presents a darker brown hue compared to the corallite walls. **Habitat:** observed in reef environments.

Montipora cf. undata Bernard, 1897



A colony characterized by its encrusting growth, featuring small, immersed, and indistinct corallites. The coenosteum exhibits tuberculae fused into ridges. Typically brown with pale growing margins. **Habitat:** observed on upper reef slopes.

Acroporidae gen. et sp. indet.



An Acroporid species with colonies forming clusters of thin, irregularly fused branches. The corallites are small and immersed, while the coenosteum is smooth. The coloration is pale brown with white tipped branches. **Habitat:** reef environments.

Leptoseris sp. indet.



St. 39. Depth: 43 m



St. 39. Depth: 41 m

Coral colonies may be horizontal to vertical with unifacial and irregular fronds, which form upright cylindrical tubes. **Habitat:** observed in reef environments.

Note: *Leptoseris* species in Indian Ocean waters typically exhibit horizontal to vertical unifacial fronds, whorls of broad fronds, or take on encrusting forms, rather than tubular structures. The species observed here could potentially represent the initial documentation of the tubular species, *Leptoseris tubulifera*, which has yet to be documented in this area.

Fungiidae sp. indet.



A solitary, free-living coral with circular and arched polyps. Tentacles are often extended during the day. **Habitat:** reef slopes or flat shallow substrates.

Fungiidae sp. indet. 1




A solitary, free-living coral with distinctly non-circular polyps. **Habitat:** reef slopes and shallow reef environments.

Turbinaria cf. mesenterina (Lamarck, 1816)



St. 39. Depth: 42 m



St. 39. Depth: 42 m

Colonies of this coral species feature unifacial laminae, which can be contorted and fuse together in subtidal environments, grow upright on upper reef slopes, or spread horizontally in deeper waters, as shown here. These growth patterns are influenced by the availability of light. Typically, colonies are less than 1 m in size but can be substantially larger in fringing reef areas. Corallites are densely packed, slightly protruding. Colour is usually grey-green or grey-brown. **Habitat:** diverse, especially in turbid waters and around reef bases.

Lobophyllia cf. hemprichii (Ehrenberg, 1834)




St. 39. Depth: 42 m



St. 39. Depth: 42 m


Colonies exhibit a flat to hemispherical morphology and can span more than 5 m in diameter. Their growth forms range from phaceloid to flabellomeandroid, with the corallites arranged in irregularly winding or maze-like patterns. Septa gradually taper in thickness from the coral wall towards the columella, featuring tall, sharp teeth. When retracted, polyps appear thick and fleshy, presenting either smooth or rough mantles. **Habitat:** reef slopes.

Astrea sp. indet.



Colonies are irregularly sub-massive or encrusting. Corallites are circular. **Habitat:** shallow environments, especially reef flats.

Dipsastraea cf. *rotumana* (Gardiner, 1899)



Often flat and subplocoid. Corallites are clearly subplocoid on convex surfaces; crowded on flat surfaces, becoming cerioid and irregularly shaped. **Habitat:** reef associated in shallow waters.

Note: the genus *Dipsastraea* was previously known as *Favia*.

Dipsastraea sp. indet.



St. 36. Depth: 25 m

Colonies are dome-shaped or flat. Corallites are thick walled and circular, ranging within and among colonies from fully plocoid to cerioid. **Habitat:** reef slopes and lagoons.

Note: the genus *Dipsastraea* was previously known as *Favia*.

Favites cf. *abdita* (Ellis & Solander, 1786)



St. 37. Depth: 33 m

Colonies are massive, either rounded or hillocky and sometimes over 1 m across. Corallites are rounded, with thick walls. **Habitat:** reef environments.

Favites cf. vasta (Klunzinger, 1879)



St. 39. Depth: 38 m

Colonies are massive and are commonly over 1 m across. Corallites are deep and angular and have very thick walls. **Habitat:** most reef environments.

Oulophyllia crispera Lamarck, 1816



St. 36. Depth: 39 m

Colonies are sub-massive or hemispherical and are frequently over 1 m across. Valleys are short, broad, V-shaped and have sharp upper margins. **Habitat:** shallow reef environments.


Paramontastraea peresi (Faure & Pichon, 1978)



St. 39. Depth: 42 m

Colonies are encrusting and helmet-shaped. **Habitat:** shallow reef environment.

Platygyra cf. acuta Veron, 2000



St. 39. Depth: 42 m

Colonies are massive and meandroid, with walls forming an acute or sharp edge. Septa are uniformly exsert and have ragged margins. **Habitat:** most reef environments, especially shallow fringing reefs.

Platygyra cf. daedalea (Ellis & Solander, 1786)



St. 39. Depth: 43 m

Colonies are generally massive, and meandroid or submeandroid, with thick walls. Septa are exsert and have a characteristically ragged appearance. **Habitat:** Most reef environments, especially back reef margins.

Family Pocilloporidae

Pocillopora sp. indet.



St. 36. Depth: 39 m

Colonies are cespitose, i.e., growing in small dense clumps, with main branches often rounded at the base. **Habitat:** shallow reef environments.

Pocillopora sp. indet. 1



Colonies are cespitose, i.e., growing in small dense clumps, with main branches often rounded at the base. **Habitat:** shallow reef environments.

Note: the photo shows a colony of *Pocillopora* sp. on the right side and a colony of *Stylophora* on the left side, identified as *S. madagascarensis* by F. Benzoni.

Stylophora sp. indet.



Colonies are ramosely, i.e., with many branches which can be slender, digitiform or palmate. **Habitat:** shallow reef environments.

Goniopora sp. indet.




St. 32. Depth: 46 m




St. 32. Depth: 46 m

Colonies are submassive. Living colonies are usually brown, either pale or dark. They are readily distinguished by having numerous daisy-like polyps that extend outward from the base, each tipped with 24 stinging tentacles which surround a mouth. **Habitat:** many reef environments including turbid waters.

At least three growth forms of *Porites* were observed in the area. Here, we distinguish between branched and massive colonies, considering the branched colonies as a separate species from the massive ones. It remains unclear whether these two growth forms represent morphological plasticity or signify distinct species.





Porites sp. indet.



Colonies of this species or morphotype exhibit knobby growth forms with short, stumpy branches. **Habitat:** reef flats.


Note: several colonies with a knobby growth form were observed at station 36. They were identified as *Porites* cf. *fontanesii* by L. De Vantier, a species that was described by Benzoni and Stefani in 2012 from the southern Red Sea and the Gulf of Aden. Here we designate it as *Porites* sp. indet. as F. Benzoni did not confirm its status as this specific species.

Porites sp. indet. 1 (Growth form I)




Colonies of this species or morphotype are cream or brown in colour, massive, reaching sizes exceeding 1 m in diameter, and exhibit a bumpy growth surface with rounded knobs. **Habitat:** reef flats.

Porites sp. indet. 2 (Growth form II)




St. 36. Depth: 26 m



St. 36. Depth: 26 m

Colonies of this species or morphotype resemble *Porites* sp. indet. 1 in their massive structure and cream or brown coloring. However, their knobs displayed distinct sharp corners. **Habitat:** reef flats.

Heliopora cf. coerulea (Pallas, 1766)



St. 36. Depth: 26 m




St. 36. Depth: 26 m

Commonly known as the blue coral due to the presence of iron salts in its skeleton. Colonies consist of thin, vertical brown branches with eight-tentacled polyps, giving the surface a hairy appearance. It has three main growth forms: columnar, lobate, and encrusting, exhibiting high morphological plasticity and continuous variation among these forms. It can form very large colonies, especially in reef lagoons. The International Union for Conservation of Nature (IUCN) classifies it as a threatened species. **Habitat:** reported from a wide range of environments, including intertidal reef flats, wave-exposed reefs, reef fronts, reef slopes (from less than 1 m to over 60 m deep), and marginal habitats.

Heliopora sp. indet.



St. 36. Depth: 39 m



St. 39. Depth: 32 m

Until 2018, *Heliopora* was known to consist of only a single valid species, *H. coerulea*, and all descriptions available actually refer to this species. However, in 2018, Richards *et al.* described a new species, *H. hiberniana*, which has a white skeleton instead of blue and is currently known only from offshore areas in northern Western Australia. The morphological descriptions of both species are very similar, except for the colour of the skeleton when broken. **Habitat:** observed on reef flats.

Family Ellisellidae

Class Octocorallia
Order Scleralcyonacea

Ellisellidae gen. et sp. indet.




St. 36. Depth: 26 m

Colonies observed are attached to a hard surface by a holdfast. They are bushy with dense branches protruding from a narrow, hollow, cross-chambered central core. The spicules are calcareous and less than 0.3 mm in length while the polyps are retractable. **Habitat:** shallow reef to mesophotic environments.

Family Isididae

| Order Malacalcyonacea

Isididae gen. et sp. indet.



St. 36. Depth: 26 m

This colony was tentatively identified as a member of the Isididae (Benayahu, pers. comm.). **Habitat:** observed on reef flats.

Rumphella sp. indet.



St. 36. Depth: 39 m



St. 39. Depth: 25 m

Possibly a species of *Rumphella* with colonies typically bushy, with either sparse, whip-like branches or dense shrub-like branches that have a smooth surface and blunt tips. Light brown to greyish colour. **Habitat:** observed on reef flats.

Lobophytum sp. indet.



St. 37. Depth: 33 m

Colonies have a thick encrusting morphology and a raised surface. They have two types of polyps: large autozooids that extend from the crown (capitulum) and small siphonozooids embedded in it. They exhibit a distinct “hairy” appearance when the autozooids are extended. *Lobophytum* possess thick-fleshed, digitate outgrowths on short stalks at the edges of the capitulum, whereas *Sarcophyton* do not. **Habitat:** shallow reef waters, but also on hard substrates in deeper waters.

Sarcophyton sp. indet.



St. 37. Depth: 34 m

Colonies have a prominent bare stalk and a large, smooth upper surface with distinct convoluted edges. As *Lobophytum*, *Sarcophyton* species have two types of polyps: large autozooids and small siphonozooids. The underside of the coral is visible where the edges are folded upwards. **Habitat:** usually found in sheltered back reef and lagoonal habitats, but in the area on hard substrates in deeper waters.

Millepora sp. indet.



Colonies have a calcareous skeleton with diverse growth-forms, from fine branching domes, encrusting sheet-like, to laminar. Colonies are brownish with a pale yellow hue. **Habitat:** observed on shallow reefs.

Note: at least three species have been reported in the Western Indian Ocean, differing in pore size and arrangement, colony branching, shape and colour. Individuals also differ depending on local conditions.

Sting is painful, burning, hence the common name, Fire Coral but not a true coral.

ECHINODERMS



Holothuria atra Jaeger, 1833



St. 39. Depth: 32 m

A sea cucumber with a uniformly black body, commonly covered with medium-grain sand, with characteristic bare circles in two rows along the dorsal surface. Podia on the dorsal surface small and sparse. Tentacles black. **Habitat:** inner and outer reef flats, and seagrass beds, 0–30 m deep. Found in Mauritius and the Saya de Malha Bank, often near *Halimeda* sp.

Holoturia coronopertusa Cherbonnier, 1980



St. 34. Depth: 46 m

A sea cucumber typically orange-brown dorsally and light-peach ventrally, with light to dark brown dorsal papillae spread over the entire surface, larger near the mouth and anus. **Habitat:** reported to forage on coralline sand, both during the day and during the night, in relatively deep waters 20–100 m deep.

Stichopus cf. pseudohorrens Cherbonnier, 1967



St. 37. Depth: 34 m

A large sea-cucumber, brownish-yellow to rosy red or brown with darker mottling. Body arched dorsally and flattened ventrally. Very long, large conical papillae dorsally. Long papillae also occur along the lateral margins of the ventral surface. The mouth is ventral, with 20 long dark brown tentacles. **Habitat:** hard bottoms with coral rubble and sand.


Thelenota ananas (Jaeger, 1833)



St. 39. Depth: 33 m

A characteristic sea cucumber with large papillae covering the dorsal surface, appearing conical, star-shaped, or somewhat branched. The colour dorsally varies from reddish-orange to brown, ventrally light pink, with podia ranging from brown to pink. Body firm, arched dorsally, flattened ventrally. Mouth ventral with 20 large, brown tentacles. **Habitat:** hard bottoms with large coral rubble, and coral patches.

Linckia cf. laevigata (Linnaeus, 1758)



St. 39. Depth: 33 m

A starfish with five cylindrical arms and a bright blue or light blue body, though other colors have been observed. It can grow up to 30–40 cm across. Identification is tentative and based on coloration; confirmation would require a close-up view of the underside. **Habitat:** shallow reef areas exposed to the sunlight, among sponges, macroalgae and seagrasses and on rocks, dead coral or rubble at depths of 0 to 60 m.

Linckia sp. indet.



St. 9. Depth: 58 m

A starfish of the genus *Linckia*, with long conical arms that are yellowish mottled with brown near the center and dark brown towards the tips. Dr. Mah tentatively identified it as *Linckia profunda*, the mesophotic species of *Linckia* described from the Pacific Ocean. **Habitat:** observed on a sand and rubble bottom at 58 m depth.


Culcita cf. novaeguineae Müller & Troschel, 1842



St. 36. Depth: 38 m

A cushion star, morphologically very similar to *C. schmideliana*, predominantly found in the eastern Indian and Pacific Oceans. Provisionally identified as *C. novaeguineae* due to the absence of large, dark granules or nodules on its body surface. **Habitat:** lagoon areas and on inner reef flats with seagrasses and among algae at depths down to about 90 m depth.


Culcita schmideliana (Retzius, 1805)



St. 39. Depth: 48 m

A peculiar cushion star with a leathery, puffed-up appearance, featuring a convex upper surface and a flat underside. Coloration varies, but large dark granules are present, with variation in their abundance. **Habitat:** lagoon areas and on inner reef flats with seagrasses and among algae at depths down to about 90 m depth.


Pentaceraster sp. indet.



St. 40. Depth: 43 m

Members of the genus *Pentaceraster* differ from those belonging to *Protoreaster*, by having spines on the marginal plates between the arms. Dr. Mah identified this specimen as possibly *P. alveolatus*, an Indo-West Pacific species commonly known as the Cushion sea star, which occurs in sandy-rocky intertidal regions and on reef platforms at depths between 1 and 60 m.

Protoreaster lincki (Blainville, 1830)



St. 39. Depth: 42 m

A large starfish, 25–30 cm, with a rigid body, white to light gray dorsally, but traversed by a vivid red network connecting large, blunt tubercles, which can be slightly sharper and denser at the ends of the arms; their shape, size, and number vary considerably. *Protoreaster* species lack spines on the marginal plates between the arms. **Habitat:** sandy bottoms and seagrass meadows from shallow waters, down to 100 m depth.


Echinothrix calamaris (Pallas, 1774)



St. 42. Depth: 49 m

Commonly known as the Black banded sea urchin, this species has two sets of spines: short, slender, yellow-to-dark spines, and longer, thicker spines, often banded. A translucent large anal sac surrounded by visual receptors for rudimentary vision is visible on the aboral side. **Habitat:** lagoons, external reef slopes and channels down to 70 m depth.

Echinothrix diadema (Linnaeus, 1758)



St. 42. Depth: 48 m

A sea urchin with long, slender spines generally black or blue-black in colour (the spines show a blue sheen in the light). Anal sac is small and dark. Their round body, or test, features five star-patterned naked zones called iridophores, which can be brightly colored. **Habitat:** coral reefs, rocky substrates, and seagrass beds, often in shallow waters but can also be found at greater depths.


Cidaridae gen. and sp. indet.



A sea urchin belonging to the family Cidaridae. Members of this family have robust, thick spines and their body, or test, is typically covered in secondary spines and tubercles, giving them a rough appearance. **Habitat:** shallow coastal habitats to the deep ocean floor.

Family Toxopneustidae | Order Camarodonta

Toxopneustes pileosus (Lamarck, 1816)



Commonly known as the Flower urchin, it is notable for its colorful, petal-like pedicellariae which can resemble flowers. It has short, blunt spines that are often hidden by these pedicellariae. This species is venomous and can deliver painful stings. **Habitat:** shallow coastal habitats.

OTHER INVERTEBRATES



Conus cf. tessulatus Born, 1778



A sea snail species, robust and conical in shape, with a highly variable coloration that generally features a pattern of tessellated or checkered markings. The base color can range from white to yellowish or pale brown, with darker brown, orange or reddish tessellated patches. The base of the shell is sometimes dark colored. **Habitat:** shallow waters, commonly found on sandy or rocky substrates, coral reefs, inner reef platforms, and deeper areas.


Family Strombidae

Lambis cf. lambis (Linnaeus, 1758)



A sea snail species with a thick and heavy shell, variable in colour, often displaying a combination of brown, cream, orange, and pink hues. The exterior of the shell may be mottled or patterned. The aperture is wide and flaring. The outer lip is flared, thickened and expanded, with six distinctive finger-like projections or spines of the shell (that resemble spider legs) radiating outward. **Habitat:** mangrove areas, seagrass beds, reef flats and coral-rubble bottoms in shallow to deeper waters.

Phyllidiopsis quadrilineata (Bergh, 1905)



St. 37. Depth: 38 m

A nudibranch species with a very light background colour and a distinct pattern of longitudinal black lines on the dorsum. The rhinophores (a pair of external chemosensory structures on the head) and rhinotubercles (tubercles on the mantle) are yellow. There are three to five, longitudinal, blue grey to pale grey ridges (broken or continuous) joining the tubercles. A central ridge is always present. Towards the margins of the dorsal surface or notum, there are usually short, transverse ridges. **Habitat:** frequently found on the fore reef zone and the reef flat zone. Some are also found in moderately deep to shallow waters.


Odontodactylus japonicus (De Haan, 1844)



St. 9. Depth: 57 m

A Mantis shrimp species overall salmon in colour. Uropod yellow; exopod with outer movable spines yellow orange with blue posterior margin; endopod and distal segment of exopod with red setae. **Habitat:** sandy and shelly bottoms.

Odontodactylus latirostris Borradaile, 1907



St. 39. Depth: 53 m

A Mantis shrimp species overall mottled light brown on white-cream, with darker brown mid-dorsal surface. Uropod with black band across proximal segment of exopod, adjacent protopod and endopod; exopod distal segment pink. **Habitat:** sandy and shelly bottoms.

BONY AND CARTILAGINOUS FISHES



Gymnothorax favagineus Bloch & Schneider, 1801



St. 36. Depth: 38 m

A moray eel with moderately sized teeth, bold coloration of narrow yellow reticulations enclosing dark brown to black polygons (dark honey-comb-like pattern); colour of uniform intensity on head, body and fins, and continuing into mouth. **Habitat:** rocks and crevices of reef flats and outer reef slopes of continental reefs. **Note:** the cardinal fish *Ostorhinchus nigrofasciatus* at the bottom right.

Myripristis botche Cuvier, 1829



St. 39. Depth: 41 m

A soldierfish with head red anteriorly; postorbital head and body silvery white, edges of scales dark; opercular membrane black from about ½ distance from opercular spine to pectoral-fin base; spinous dorsal fin whitish basally; other median fins red with white leading edges, and lobe tips with large black spots; paired fins whitish. **Habitat:** reef-associated.

***Myripristis* sp. indet.**



St. 37. Depth: 37 m

A soldierfish with red body, large scales with darker edges; black on opercular membrane; fins with white leading edge (except for pectoral fins). **Habitat:** reef-associated.

Sargocentron* cf. *seychellense (Smith & Smith 1963)



St. 37. Depth: 37 m

A squirrelfish with alternating stripes of brownish red (darker dorsally) on body; 2 pairs of brownish red stripes on sides of body converging on peduncle; head red dorsally, yellowish ventrally; opercle silvery with 2 brownish red bars, and an oblique reddish brown band from eye to corner of preopercle; pectoral-fin base red, axils dark brown; spinous dorsal fin with pale red spines, other fins mainly yellowish; upper and lower margins of caudal fin red. **Habitat:** shallow waters, coral reefs and rocky shores.


Sargocentron cf. diadema (Lacepède 1801)



St. 39. Depth: 32 m

A squirrelfish with alternating stripes of red and silvery white on body, red stripes much broader than white ones; spinous dorsal fin deep red to reddish black, membrane tips white, usually with curving longitudinal white band along lower third of fin. **Habitat:** coral reefs, between 2 and 42 m depth.

Apogonichthyoides pharaonis (Bellotti, 1874)



St. 40. Depth: 49 m

Cardinal fish with body generally bronzy dorsally and silvery ventrally, with 3 dark brown to black bars, 1st bar joining dark leading edge of 1st dorsal fin and extends downwards through large ocellated dark spot and fading near ventrum; 2nd bar extending from middle of 2nd dorsal fin to middle of anal fin; 3rd bar across rear of peduncle; cheeks with narrow dark mark from eye to preopercle ridge. **Habitat:** silty reefs, mangroves, seagrass beds..


Amblyeleotris periophthalma (Bleeker, 1853)



A shrimpgoby with head and body whitish to pale yellowish, 5 orange to reddish brown blotchy bands; pale interspaces on body with brownish to orange mottling and spots; similarly coloured round spots on head; 2 red spots or 1 blotch at corners of mouth; caudal-fin base with a curved brownish bar.

Habitat: sandy and coral-rubble substrates, between 10 and 25 m depth.

Valenciennesa cf. wardii (Playfair, 1867)



St. 9. Depth: 57 m




St. 39. Depth: 53 m

A glidergoby with 1st dorsal fin rounded; head and body pale grey to pearly white; 3 brown dark-margined bars on body and similar bar across distal half of caudal fin; oblique iridescent blue-white stripes below eyes, on sides of head; rear of 1st dorsal fin with large black spot, and 2 black bars on lower part of 2nd dorsal fin.

Habitat: silty sand, observed between 16 to 60 m depth.

Family Trichonotidae

Trichonotus cf. marleyi (Smith, 1936)



St. 9. Depth: 56 m

A sand-diver with a yellowish body dorsally, pale reddish ventrally with translucent bluish green and red spots; iris flap golden. Males with up to 14 indistinct dark markings and 3 longitudinal rows of small brilliant bluish white spots on upper half of body; fins transparent; dorsal- and anal-fin rays with small yellowish dots. Females with indistinct narrow dark stripe above lateral line; dorsal and caudal fins with small yellow to dark brown spots. **Habitat:** sandy bottoms.

Family Callionymidae

Class Actinopterygii
Order Sygnathiformes


Synchiropus cf. sechellensis Regan, 1908



A species of dragonets with snout shorter than eye diameter, head and body rose-pink or red-orange, white ventrally, with irregular broad red bars and narrower white blotches; 1st dorsal fin of males with small brown ocelli, fin of females alternatively spotted with black and white; caudal fin with 2 red-orange bars. **Habitat:** soft bottoms, 37 to 68 m depth.

Family Fistulariidae

Fistularia cf. commersonii Rüppell, 1838



A flutemouth species with a body uniformly olive-green or brownish dorsally, with pair of blue or green lines and/or rows of blue spots; silvery white ventrally; median fins pinkish distally; caudal filament white. **Habitat:** coral reefs and seagrass beds, to at least 120 m depth.

Carangidae gen. et sp. indet.



St. 37. Depth: 36 m

A trevally or scad with fusiform body, compressed laterally. Dorsal fin usually divided into separate spinous and soft-rayed fins; caudal fin strong, widely forked, with slender peduncle; scales in posterior (straight) part of lateral line often modified as scutes. **Habitat:** pelagic, usually in schools.

Family Echeneidae

Echeneis naucrates Linnaeus, 1758



St. 34. Depth: 46 m

A remora with robust, elongated body; sucking disc large, with 21–28 laminae; caudal fins convex to almost truncate in adults, and lanceolate in juveniles. Body brown to black, with pale bands. **Habitat:** near and far from the coast, often free-swimming in shallow inshore areas and around coral reefs. Temporarily attaches to hosts like sharks, rays, large fish, sea turtles, whales, dolphins, and ships.

Variola louti (Forsskål, 1775)



A lyretail with variable colour: head, body, and median fins range from yellowish-brown to orange-red, covered with small blue, lavender, or pink spots. Median-fin margins are yellow, pectoral fins red to brown with yellow tips. **Habitat:** offshore coral reefs, usually > 15 m depth, in clear waters.

Remarks: in Mauritius, a specimen caught from oceanic banks weighing more than 3 kg when ungutted or more than 2.6 kg when gutted is considered toxic. If caught in Mauritius's territorial waters, it is considered toxic if it weighs more than 1.5 kg when ungutted or more than 1.3 kg when gutted.

Family Scorpaenidae

Dendrochirus hemprichi Matsunuma, Motomura & Bogorodsky, 2017



A species of lionfish with dark reddish brown to scarlet red body, with 5 or 6 broad darker saddles, sometimes with pale areas between them; 3 indistinct dark red bars below eyes; dorsal fin spines red or reddish brown, with small dark spots; pectoral fins with 4–10 dark brown to reddish bars, alternating with pale whitish to yellowish area. **Habitat:** reef-associated, to about 70 m depth.

Pterois miles (Bennett, 1828)



St. 39. Depth: 47 m

A firefish with numerous dark red or brown vertical bars, of irregular width; 4 or 5 stripes radiate from the eyes; supraocular cirrus usually dark brown or black; dorsal-fin spines with dark brown, dark grey or dark red areas alternating with nearly white areas; dorsal, anal and caudal fin rays with numerous black spots, membranes translucent. **Habitat:** reef-associated.

Family Zanclidae

Zanclus cornutus (Linnaeus, 1758)



St. 39. Depth: 30 m

Known as the Moorish idol, it has compressed disc-like body, with 2 broad vertical black bars (one bar from dorsal fin origin, across head and covering anterior body and pectoral and pelvic fins, the 2nd bar on rear of body; caudal fin black with white margin; dorsal fin extending into a long filament. **Habitat:** coral and rocky reefs.

Naso brevirostris (Cuvier, 1829)



St. 37. Depth: 33 m

A unicorn fish with a broad, tapering horn extending up to one head length forward of the mouth in adults, appearing as a bump in fish under 10 cm total length; moderately developed keels on each side of the peduncle; caudal fin truncate to slightly rounded. Body pale bluish-grey to olive-brown with fine vertical dark grey lines on midsides, ending as rows of spots ventrally; head with small dark spots or reticular lines, and horn with oblique dark lines. **Habitat:** reef-associated.

Naso hexacanthus (Bleeker 1855)



St. 39. Depth: 34 m

A brownish to bluish-grey surgeonfish becoming yellowish below, with a black margin on the cheek and gill cover. Capable of rapidly changing to pale greyish blue overall, as when at a wrasse cleaning station. **Habitat:** inshore reefs and coral-rich areas.

Diagramma centurio Cuvier, 1830



St. 37. Depth: 38 m

A species of rubberlip with dusky greyish body, with numerous small brownish yellow spots on head, back, sides, upper part of caudal fin, and sometimes over entire caudal fins. **Habitat:** reef-associated.

Family Lutjanidae

Lutjanus sebae (Cuvier, 1816)



St. 39. Depth: 30 m

A snapper with reddish body, becoming paler ventrally. Juveniles and subadults silvery white, with 3 broad reddish brown bands (1st from snout tip to dorsal-fin origin, 2nd vertical through spinous dorsal fin to pelvic fins, 3rd from beneath anterior dorsal-fin rays to lower leading edge of caudal fin); bars fade with growth and body become redder. **Habitat:** coral or rocky reefs, to about 180 m depth.

Lethrinus enigmaticus Smith, 1959



A greyish or tan emperor, with grey or brownish head. Dark purplish spots around front and lower rim of eyes; pale yellowish streak radiating forward from eyes to nostrils, and yellowish cross-stripes on upper head; upper sides often with about 7 dark bars, sometimes 3 bronze stripes ventrally. **Habitat:** seagrass beds, coral reefs and adjacent sandy areas, usually to about 50 m depth.

Lethrinus variegatus Valenciennes, 1830



A slender grey or brown emperor, sometimes with scattered irregular dark blotches on sides; often 2 dark bands below eyes; inner surface of pectoral fins never red in life. **Habitat:** sandy areas near coral reefs, to about 150 m depth.


Family Nemipteridae

Scolopsis frenata (Cuvier, 1830)



A species of monocle bream with body blue or olive-green on dorsum, white below; snout dusky; narrow blue stripe from snout tip to anteroventral part edge of eye, yellow stripe from top of snout, through upper part of eyes, arching on back to upper part of peduncle; stripe edged dark green above from behind eyes to beneath dorsal fin. **Habitat:** sandy bottoms near coral reefs.

Pomacanthus imperator Bloch, 1784



St. 36. Depth: 40 m

A characteristic angel fish with alternating yellow and blue stripes on body extending onto dorsal and anal fins; dark mask edged with blue over eyes; nape yellowish or greenish grey; snout and cheek bluish white; caudal fin yellow.

Habitat: lagoons and outer-reef slopes, in 1 to 60 m depth.

Pomacanthus semicirculatus (Cuvier 1831)



St. 39. Depth: 42 m

An angel fish with head free of spots, body yellowish grey that transitions to a dusky brown towards the distal areas, adorned with small, dark blue oval spots. This pattern continues onto the median fins, where the spots become rounder; fin margins are bright blue. **Habitat:** inshore reefs and coral-rich areas.

Oxycheilinus bimaculatus (Valenciennes, 1840)



St. 36. Depth: 38 m

A wrasse with colour variations, females reddish-orange to orangish-brown hues, marked with whitish blotches and yellow ventrally, and a prominent blackish blotch between the pectoral-fin tips and lateral line. Males orangish-brown and pale green, featuring a conspicuous blackish blotch on the sides. Head is predominantly green with radiating pink to orange-red lines from the eyes. **Habitat:** areas with heavy growth of algae or seagrass, to 100 m depth.


Chlorurus sp. indet.



St. 36. Depth: 26 m

A parrotfish with moderately deep body; teeth in jaws fully fuse to form dental plates which are fully exposed, outer surface smooth and cutting edge irregular. **Habitat:** reef-associated.

Sufflamen chrysopterum (Bloch & Schneider, 1801)



St. 37. Depth: 40 m

A dark brown triggerfish with bluish purple throat and belly; short transverse white band on chin; vertical band varying in colour from yellow to pale brown, extending from back edge of eye to lower pectoral-fin base; caudal fin dark brown with broad white rear margin and narrow upper and lower white margins.

Habitat: shallow lagoons and sheltered back reefs, to about 20 m depth.

Family Ostraciidae

Lactoria fornasini (Bianconi, 1846)



St. 36. Depth: 40 m


A cowfish with distinct mid-dorsal ridge and stout retrorse spine. Body enclosed in hard armoured-like carapace with 5 ridges, cross-section pentagonal in adults. Carapace comprises pale brown or yellowish scale plates, with central blue or magenta spot that can change colour within seconds. Spines ('horns') present in front of eyes. **Habitat:** coral reefs and weedy areas, at 5 to 80 m depth.

Arothron hispidus (Linnaeus 1758)



A pufferfish with head and body greenish brown dorsally, with many small white spots, becoming white ventrally, with curved dark stripes; wide dark vertical bars on sides of head and body; pectoral-fin bases and gill opening encircled with alternating white or yellow and black lines. **Habitat:** coral reefs, to about 59 m depth. Juveniles usually occur in weedy areas of estuaries.

Canthigaster rivulata (Temminck & Schlegel, 1850)



Head and body brownish grey dorsally, with many short, wavy, dark blue lines creating vermiculate pattern; lower half of body white; sides with 2 parallel dark bands, joined by vertical section curving around front of gill opening; dark blotch at dorsal-fin base; dorsal, anal and pectoral fins pale; caudal fin with yellowish brown rays and dark mark at base of lower rays. Attains 20 cm total length. **Habitat:** rocky and coral reefs.

Taeniurops meyeri (Müller & Henle 1841)



A stingray with a nearly circular disc, slightly wider than long, all outer margins evenly rounded, relatively short tail, less than disc width, with usually 1 stinging spine, and small pelvic fins small. Dorsal surface of disc usually dark purplish, bluish or brownish grey dorsally, or pale with scattered irregular dark spots or mottling and dusky specks or streaks; disc whitish ventrally. **Habitat:** bottom-dwelling in lagoons, estuaries and on reefs.

REFERENCES

- Bergstad, O.A., Bissessur, D., Sauba, K., Rama, J., Coopen, P., Oozeeraully, Y., Seeboruth, S., Audit-Manna, A., Nicolas, A., Reetoo, N., Tabachnick, K., Kuyper, D., Gendron, G., Hollanda, S., Melanie, R., Souffre, A., Harlay, J., Bhagooli, R., Soondur, M., Ramah, S., Caussy, L., Ensrud, T.M., Olsen, M. and Høines, Å.S. 2018. *Regional Resources and Ecosystem Survey in the Indian Ocean, Leg 2.1: Characterizing Ecosystems and Morphology of the Saya de Malha Bank and Nazareth Bank, 3 May - 4 June 2018*. NORAD-FAO Programme GCP/GLO/690/NOR, Cruise Reports *Dr. Fridtjof Nansen*, EAF-Nansen/CR/2018/6.
- Hagan, A. and Robinson, J. 2001. Benthic habitats of the Saya de Malha Bank. Marine Science, training, and education in the western Indian Ocean. *Royal Geographical Society, London/30 Reefs Symposium of the Zoological Society of London*, 28: 26–27.
- Hilbertz, W., Gutzeit, F. and Goreau, T. 2002. *Saya de Malha Expedition March 2002*, funded by the Lighthouse Foundation, pp. 1–107.
- Kohler, K.E., and Gill, S.M. 2006. Coral Point Count with Excel extensions (CPCe): A Visual Basic program for the determination of coral and substrate coverage using random point count methodology. *Computers & geosciences*, 32(9), 1259–1269.
- Obura, D.O., Church, J.E. and Gabrié, C. 2012. *Assessing marine world heritage from an ecosystem perspective: the Western Indian Ocean*. World Heritage Centre, United Nations Education, Science and Cultural Organization (UNESCO). 124 pp.
- Richards, Z.T., Yasuda, N., Kikuchi, T., Foster, T., Mitsuyuki, C., Stat, M., Suyama, Y. and Wilson, N.G. 2018. Integrated evidence reveals a new species in the ancient blue coral genus *Heliopora* (Octocorallia). *Scientific Reports*, 8(1), 15875. <https://doi.org/10.1038/s41598-018-32969-z>.
- Rosen, B. 1971. The distribution of reef coral genera in the Indian Ocean. Regional variation in Indian Ocean coral reefs. *Symposium of the Zoological Society of London*, 28: 263–299.
- Serigstad, B., Ensrud, T.M., Olsen, M., Ostrowski, M., Adu Kumi, S., Akoto, L., Aggrey-Fynn, J., Blazewicz-Pasz-kowycz, M., Jozwiak, P., Pabis, K., Siciński, J. 2015. *Environmental monitoring Ghana 2012, chemical and biological analysis*. NORAD – FAO Project GCP/003/NOR Survey Report *R/V Dr Fridtjof Nansen* EAF-Nansen/2012/7. 163 pp.
- Webber, T., Lewis, T., Talma, S., Sandoo, S., Meller, L., Dulau-Drouot, V., and Thompson, K.F. 2023. *Cetaceans of the Saya de Malha Bank region, Indian Ocean: A candidate Important Marine Mammal Area*. Regional Studies in Marine Science. <https://doi.org/10.1016/j.rsma.2023.103164>.

FURTHER READINGS

Bhagooli, R., Ramah, S., Kaullysing, D., Gopeechund, A. and Bergstad, O.A. 2021. First field observations of *Halimeda* beds at depths of 37–62 m at Saya de Malha and Nazareth Banks, Mascarene Plateau. *WIO Journal of Marine Science*, Special Issue, 2/2021, 183–189.

Bhagooli, R. and Kaullysing, D. 2019. Seas of Mauritius. In C. Sheppard (Ed.), *World Seas: An Environmental Evaluation* (2nd ed., Vol. II: The Indian Ocean to the Pacific). Academic Press. ISBN: 9780081008539.

New, A. L., Standfield, K., Smythe-Wright, D., Smeed, D.A., Evans, K. and Alderson, S.G. 2005. Physical and biochemical aspects of the flow across the Mascarene Plateau in the Indian Ocean. *Philosophical Transactions of the Royal Society A*, 363, 151–168. <https://doi.org/10.1098/rsta.2004.1484>.

Ramah, R. and Bhagooli, R. 2021. First field observation of a *Thalassodendron ciliatum* bed on the Nazareth Bank, Mascarene Plateau. *WIO Journal of Marine Science*, Special Issue, 2/2021 179–182.

Ramah, S., Kaullysing, D. and Bhagooli, R. 2021. Sightings of sea stars (Echinodermata, Asteroidea) and a first record of crown-of-thorns starfish *Acanthaster* at Saya de Malha Bank, Mascarene Plateau. *WIO Journal of Marine Science*, Special Issue, 2/ 2021, 197–201.

Ramah, S., Bhagooli, R., Kaullysing, D. and Bergstad, O.A. 2021. Rhodolith beds (Corallinaceae, Rhodophyta): An important marine ecosystem of the Saya de Malha and Nazareth Banks. *WIO Journal of Marine Science*, Special Issue, 2/2021, 171–178.

Ramah, R., Gendron, G., Bhagooli, R., Soondur, M., Souffre, A., Melanie, R., Coopen, P., Caussy, L., Bissessur, D. and Bergstad, O.A. 2021. Diversity and distribution of the shallow water (23–50 m) benthic habitats on the Saya de Malha Bank, Mascarene Plateau. *WIO Journal of Marine Science*, Special Issue 2, 2021, 69–80.

Vortsepneva, E. 2008. *Saya de Malha Bank – An Invisible Island in the Indian Ocean: Geomorphology, Oceanology, Biology*. Lighthouse Foundation. 44 pp.

INDEX

A

Acanthuridae	84
Acanthuriformes	84-88
<i>Aciculites tulearensis</i>	27
<i>Acropora</i> sp. indet.	31
<i>Acropora</i> sp. indet. 1	32
<i>Acropora</i> sp. indet. 2	32
<i>Acropora</i> sp. indet. 3	33
Acroporidae	31-36
Acroporidae gen. and sp. indet.	36
Actinopterygii	75-91
Agariciidae	37
Agelasida	19
Agelasidae	19
<i>Agelas oxeata</i>	19
Alismatales	11, 12
<i>Amblyeleotris periphthalma</i>	78
<i>Anacropora</i> sp. indet.	34
Angel fish	88
Anguilliformes	75
Anthoathecata	56
<i>Apogonichthyoides pharaonis</i>	77
Apogonidae	77
<i>Arothron hispidus</i>	91
Asteroidea	61-63
<i>Astrea</i> sp. indet.	41
<i>Astreopora</i> sp. indet.	35
<i>Astreopora</i> sp. indet. 1	35
<i>Axinella weltnerii</i>	19
<i>Axinellida</i>	19, 20
Axinellidae	19, 20

B

Balistidae	90
Beryciformes	75-77
Black banded sea urchin	64
Blue coral	51
BONY AND CARTILAGINOUS FISHES	73
Bryopsidales	13-15
Bubarida	21
Bubaridae	21

C

Callionymidae	80
Camarodonta	65
<i>Canthigaster rivulata</i>	91

Carangidae	81
Carangidae gen. and sp. indet.	81
Carangiformes	81
Cardinal fish	77
Caulerpaceae	13
<i>Caulerpa cupressoides</i>	13
<i>Caulerpa taxifolia</i>	13
Chalinidae	24
Chalinidae gen. et sp. indet.	24
<i>Chlorurus</i> sp. indet.	89
Cidaridae	65
Cidaridae gen. and sp. indet.	65
Cidaroida	65
Clioanida	22
Clioanidae	22
Conidae	69
<i>Conus</i> cf. <i>tessulatus</i>	69
CORALS	29
Cowfish	90
Crustacea	71
<i>Culcita</i> cf. <i>novaeguineae</i>	62
<i>Culcita schmideliana</i>	62
Cushion star	62
Cymodoceaceae	11

D

Dasyatidae	92
Demospongiae	19-27
<i>Dendrochirus hemprichi</i>	82
Dendrophylliidae	39
Diadematidae	64
Diadematoida	64
Dictyoceratida	23
<i>Dipsastraea</i> cf. <i>rotumana</i>	41
<i>Dipsastraea</i> sp. indet.	42
<i>Dragmacidon coccineum</i>	20
Dragonets	80

E

Echeneidae	81
<i>Echeneis naucrates</i>	81
ECHINODERMS	57
Echinoidea	64, 65
<i>Echinothrix calamaris</i>	64
<i>Echinothrix diadema</i>	64
Elasmobranchii	92
Ellisellidae	53

Ellisellidae gen. and sp. indet.	53
Emperor fish	86, 87
Epinephelidae	82

F

<i>Favites cf. abdita</i>	42
<i>Favites cf. vasta</i>	43
Fire Coral	56
Firefish	83
<i>Fistularia cf. commersonii</i>	80
Fistulariidae	80
Flower urchin	65
Flutemouth	80
Fungiidae	38
Fungiidae sp. indet.	38
Fungiidae sp. indet. 1	38

G

Gastropoda	69
Glidergoby	79
Gobiidae	78, 79
Gobiiformes	77-79
<i>Goniopora</i> sp. indet.	47
<i>Gymnothorax favagineus</i>	75

H

Haemulidae	85
<i>Haliclona</i> sp. indet.	24
Halimedaceae	14
<i>Halimeda</i> sp. indet.	14
<i>Halimeda</i> sp. indet. 1	14
<i>Halophila decipiens</i>	12
Haplosclerida	24, 25
<i>Heliopora cf. coerulea</i>	51
<i>Heliopora hiberniana</i>	52
<i>Heliopora</i> sp. indet.	52
Helioporidae	51, 52
<i>Hemiasterella vasiformis</i>	26
Hemiasterellidae	26
Hexacorallia	31-50
<i>Higginsia pulcherrima</i>	20
Holocentridae	75-77
<i>Holothuria atra</i>	59
Holothuriida	59, 60
Holothuriidae	59
Holothuroidea	59, 60
<i>Holoturia coronopertusa</i>	59
Hydrocharitaceae	12
Hydrozoa	56

<i>Hyrtios erectus</i>	23
------------------------	----

I

Isididae	53, 54
Isididae gen. and sp. indet.	53

L

Labridae	89
Labriformes	89
<i>Lactoria fornasini</i>	90
<i>Lambis cf. lambis</i>	69
<i>Leptoseris</i> sp. indet.	37
Lethrinidae	86, 87
<i>Lethrinus enigmaticus</i>	86
<i>Lethrinus variegatus</i>	87
<i>Linckia cf. laevigata</i>	61
<i>Linckia profunda</i>	61
<i>Linckia</i> sp. indet.	61
Lionfish	82
<i>Liosina paradoxa</i>	26
<i>Lobophyllia cf. hemprichii</i>	40
Lobophylliidae	40
Lutjanidae	85
<i>Lutjanus sebae</i>	85
Lyretail	82

M

Magnoliopsida	11, 12
Malacalcyonacea	53-55
Mantis shrimp	71
Merulinidae	41-45
<i>Millepora</i> sp. indet.	56
Milleporidae	56
Mollusca	69, 70
Monocle bream	87
<i>Montipora cf. undata</i>	36
Moorish idol	83
Moray eel	75
Muraenidae	75
Myliobatiformes	92
<i>Myripristis botche</i>	75
<i>Myripristis</i> sp. indet.	76

N

<i>Naso brevirostris</i>	84
<i>Naso hexacanthus</i>	84
Nemipteridae	87
Nudibranchs	70

Nudibranchia 70

O

Octocorallia 51-55
Odontodactylidae 71
Odontodactylus japonicus. 71
Odontodactylus latirostris. 71
Ophiasteridae 61, 63
Oreasteridae 62
Ostraciidae 90
OTHER INVERTEBRATES 67
Oulophyllia crispa 43
Oxycheilinus bimaculatus 89

P

Paramontastraea peresi. 44
Parrotfish 89
Pentaceraster alveolatus 63
Pentaceraster sp. indet. 63
Perciformes 82, 83
Petrosiidae 25
Phakellia sp. indet. 21
Phyllidiidae 70
Phyllidiopsis quadrilineata. 70
Platygyra cf. *acuta* 44
Platygyra cf. *daedalea* 45
Pocillopora sp. indet. 45
Pocillopora sp. indet. 1 46
Pocilloporidae 45, 46
Pomacanthidae 88
Pomacanthus imperator 88
Pomacanthus semicirculatus. 88
Porites sp. indet. 48
Porites sp. indet. 1 (Growth form I) 49
Porites sp. indet. 2 (Growth form II) 50
Poritidae 47-50
Protoreaster lincki 63
Pterois miles 83
Pufferfish 91

R

Remora 81
Rubberlip 85
Rumphella sp. indet. 54

S

Sand-diver 79
Sarcophytidae 55

Sarcophyton sp. indet. 55
Sargocentron cf. *seychellense* 76
Scad 81
Scleractinia 31-50
Scleralcyonacea 51-53
Scleritodermidae 27
Scolopsis frenata 87
Scopalinida 21
Scopalinidae 21
Scorpaenidae 82, 83

SEAGRASSES AND SEAWEEDS 9

Sea snails 69
Shrimpgoby 78
Snapper fish 85
Soldierfish 75, 76
Spheciospongia inconstans 22

SPONGES 17

Spongia sp. indet. 23
Spongiidae 23
Squirrelfish 76, 77
Stelligeridae 20
Stichopodidae 60
Stichopus cf. *pseudohorrens* 60
Stingray 92
Stomatopoda 71
Strombidae 69
Stylissa carteri 21
Stylophora sp. indet. 46
Sufflamen chrysopterum 90
Surgeonfish 84
Sygnathiformes 80
Synchiropus cf. *sechellensis* 80

T

Taeniurops meyeri. 92
Tethyida 26
Tetracninelida 27
Tetraodontidae 91
Tetraodontiformes 90, 91
Thalassodendron ciliatum 11
Thelenota ananas 60
Thorectidae 23
Toxopneustes pileosus 65
Toxopneustidae 65
Trevally 81
Trichonotidae 79
Trichonotus cf. *marleyi*. 79
triggerfish 90
Turbinaria cf. *mesenterina* 39

U

Udoteaceae	15
<i>Udotea</i> sp. indet.	15
Ulvophyceae	13-15
Unicorn fish	84

V

<i>Valenciennea</i> cf. <i>wardii</i>	79
Valvatida.	61-63
<i>Variola louti</i>	82

W

Wrasse	89
--------------	----

X

<i>Xestospongia testudinaria</i>	25
--	----

Z

Zanclidae	83
<i>Zanclus cornutus</i>	83



This photographic catalogue presents an overview of the flora and fauna observed in the shallow waters (less than 100 m depth) of the Saya de Malha Bank in the Mascarene Plateau. While not a comprehensive representation, as some taxa were excluded due to poor-quality photos that could result in misleading or unreliable depictions, it encompasses a total of 113 taxa identified at the family, genus, or species level. These taxa span key groups, including seaweeds, seagrasses, sponges, corals, echinoderms, crustaceans, molluscs, and fish.

The images are snapshots derived from video observations captured during the 2018 R/V *Dr. Fridtjof Nansen* ecosystem survey, which aimed to characterize the marine ecosystem and morphology of the Saya de Malha Bank. This area is notable for its rich marine biodiversity, supporting a diverse array of species and habitats that contribute to the overall health of oceanic environments.

This catalogue serves as a valuable resource for scientists conducting further explorations in the Mascarene Plateau region, providing a basic tool for taxonomic identification. By raising awareness of the significance of the Saya de Malha Bank, we aim to foster a greater appreciation for its unique marine life and promote the conservation of these critical habitats.

ISBN 978-92-5-139479-3



9 789251 394793

CD3735EN/1/12.24